

On entropic, distributional, and differential pathways

A. M. Mathai^{1,2*} and H. J. Haubold^{3**}

¹*Department of Mathematics and Statistics, McGill University, Montreal, Canada H3A 2K6*

²*Centre for Mathematical Sciences, Pala Campus, Arunapuram P.O., Pala 686 574, India*

³*Office for Outer Space Affairs, United Nations, Vienna International Centre,
P.O. Box 500, A-1400 Vienna, Austria*

*Dedicated to the Memory of Prof. Dr.habil. Dr.h.c.mult. Hans-Juergen Treder (1928-2006),
former Director of the Einstein Laboratory for Theoretical Physics, Caputh[†]*

Abstract. Product probability property, known in the literature as statistical independence, is examined first. Then generalized entropies are introduced, all of which give generalizations to Shannon entropy. It is shown that the nature of the recursivity postulate automatically determines the logarithmic functional form for Shannon entropy. Due to the logarithmic nature, Shannon entropy naturally gives rise to additivity, when applied to situations having product probability property. It is argued that the natural process is non-additivity even in product probability property situations and additivity can hold due to the involvement of a recursivity postulate leading to a logarithmic function. Generalized entropies are introduced and some of their properties are examined. Particularly, situations are examined where a generalized entropy of order α leads to entropic pathway models, exhibiting exponential and power law behavior. Subsequently it is shown that these models link to distributional and differential pathways. Connection of the generalized entropy of order α to Kerridge's measure to create "inaccuracy" is also explored. Further for each of the three pathways their relevance to Tsallis statistics and Beck-Cohen superstatistics is emphasized.

Keywords : pathway model – generalized entropy measures – Boltzmann-Gibbs entropy – Tsallis entropy

*e-mail: *mathai@math.mcgill.ca **hans.haubold@unvienna.org

[†]Commemorating an anniversary of research work see S. Antoci and D.-E. Liebscher: Hans-Juergen Treder and the discovery of confinement in Einstein's unified field theory, arXiv:0706.3989v1.

1. Entropic pathway

Mathai & Rathie (1975) consider a number of generalizations of Shannon entropy (Shannon 1948), called entropies of order α , and give various properties, including additivity property, and characterization theorems. Recently, Mathai & Haubold (2006, 2006a) explored a generalized entropy of order α , which is connected to a measure of uncertainty in a probability scheme, Kerridge's (Kerridge 1961) concept of inaccuracy in a scheme, and pathway models that are considered in this paper. It is shown that such a generalized entropy can be used to explore pathways of entropies, respective distribution functions, and related differential equations.

As defined in Mathai & Haubold (2006, 2006a) the entropy $M_{k,\alpha}(P)$ is a non-additive entropy and his measure $M_{k,\alpha}^*(P)$ is an additive entropy. It is also shown that maximization of the continuous analogue of $M_{k,\alpha}(P)$, denoted by $M\alpha(f)$, gives rise to various functional forms for f , depending upon the types of constraints on f .

Occasionally, emphasis is placed on the fact that Shannon entropy satisfies the additivity property, leading to extensivity. It will be shown that when the product probability property (PPP) holds then a logarithmic function can give a sum and a logarithmic function enters into Shannon entropy due to the assumption introduced through a certain type of recursivity postulate. The concept of statistical independence will be examined in Section 1 to illustrate that simply because of PPP one need not expect additivity to hold or that one should not expect this PPP should lead to extensivity. The types of non-extensivity, associated with a number of generalized entropies, are pointed out even when PPP holds. The nature of non-extensivity that can be expected from a multivariate distribution, when PPP holds or when there is statistical independence of the random variables, is illustrated by taking a trivariate case.

Maximum entropy principle is examined in Section 2. It is shown that optimization of measures of entropies, in the continuous populations, under selected constraints, leads to various types of models. It is shown that the generalized entropy of order α is a convenient one to obtain various probability models.

Section 3 examines the types of differential equations satisfied by the various special cases of the distributional and thus entropic pathway model.

1.1 Product probability property (PPP) or statistical independence of events

Let $P(A)$ denote the probability of the event A . If the definition $P(A \cap B) = P(A)P(B)$ is taken as the definition of independence of the events A and B then any event $A \in S$, and S the sure event are independent. But A is contained in S and then the definition of independence becomes inconsistent with the common man's vision of independence.

Even if the trivial cases of the sure event S and the impossible event ϕ are deleted, still this definition becomes a resultant of some properties of positive numbers. Consider a sample space of n distinct elementary events. If symmetry in the outcomes is assumed then we will assign equal probabilities $\frac{1}{n}$ each to the elementary events. Let $C = A \cap B$. If A and B are independent then $P(C) = P(A)P(B)$. Let

$$P(A) = \frac{x}{n}, P(B) = \frac{y}{n}, P(C) = \frac{z}{n}.$$

Then

$$\binom{x}{n} \binom{y}{n} = \binom{z}{n} \Rightarrow nz = xy, x, y, z = 1, 2, \dots, n - 1, z < x, y \tag{1}$$

deleting S and ϕ . There is no solution for x, y, z for a large number of n , for example, $n = 3, 5, 7$. This means that there are no independent events in such cases and it sounds strange from a common man's point of view.

The term "independence" of events is a misnomer. This property should have been called product probability property or PPP of events. There is no reason to expect the information or entropy in a joint distribution to be the sum of the information contents of the marginal distributions when the PPP holds for the distributions, that is when the joint density or probability function is a product of the marginal densities or probability functions. We may expect a term due to the product probability to enter into the expression for the entropy in the joint distribution in such cases. But if the information or entropy is defined in terms of a logarithm, then naturally, logarithm of a product being the sum of logarithms, we can expect a sum coming in such situations. This is not due to independence or due to the PPP of the densities but due to the fact that a functional involving logarithm is taken thereby a product has become a sum. Hence not too much importance should be put on whether or not the entropy on the joint distribution becomes sum of the entropies on marginal distributions or additivity property when PPP holds.

1.2 How is logarithm coming in Shannon's entropy?

Several characterization theorems for Shannon entropy and its various generalizations are given in Mathai & Rathie (1975). Modified and refined versions of Shannon's own postulates are given as postulates for the first theorem characterizing Shannon entropy in Mathai & Rathie (1975). Apart from continuity, symmetry, zero-indifference and normalization postulates the main postulate in the theorem is a recursivity postulate, which in essence says that when the PPP holds then the entropy will be a weighted sum of the entropies, thus in effect, assuming a logarithmic functional form. The crucial postulate is stated here. Consider a multinomial population $P = (p_1, \dots, p_m)$, $p_i > 0$, $i = 1, \dots, m$, $p_1 + \dots + p_m = 1$, that is, $p_i = P(A_i)$, $i = 1, \dots, m$, $A_1 \cup \dots \cup A_m = S$, $A_i \cap A_j = \phi$, $i \neq j$. If any p_i can take a zero value also then zero-indifferent postulate, namely that the entropy remains the same when an impossible event is incorporated into the scheme, is to be added. Let $H_n(p_1, \dots, p_n)$ denote the entropy to be defined. Then the crucial recursivity

postulate says that

$$\begin{aligned} & H_n(p_1, \dots, p_{m-1}, p_m q_1, \dots, p_m q_{n-m+1}) \\ &= H_m(p_1, \dots, p_m) + p_m H_{n-m+1}(q_1, \dots, q_{n-m+1}) \end{aligned} \quad (2)$$

$\sum_{i=1}^m p_i = 1$, $\sum_{i=1}^{n-m+1} q_i = 1$. This says that if the m -th event A_m is partitioned into independent events $P(A_m \cap B_j) = P(A_m)P(B_j) = p_m q_j$, $j = 1, \dots, n - m + 1$ so that $p_m = p_m q_1 + \dots + p_m q_{n-m+1}$ then the entropy $H_n(\cdot)$ becomes a weighted sum. Naturally, the result will be a logarithmic function for the measure of entropy.

There are several modifications to this crucial recursivity postulate. One suggested by Tverberg is that $n - m + 1 = 2$ and $q_1 = q, q_2 = 1 - q$, $0 < q < 1$ and $H_2(q, 1 - q)$ is assumed to be Lebesgue integrable in $0 \leq q \leq 1$. Again a characterization of Shannon entropy is obtained. In all the characterization theorems for Shannon entropy this recursivity property enters in one form or the other as a postulate, which in effect implies a logarithmic form for the entropy measure. Shannon entropy S_k has the following form:

$$S_k = -A \sum_{i=1}^k p_i \ln p_i, \quad p_i > 0, \quad i = 1, \dots, k, \quad p_1 + \dots + p_k = 1, \quad (3)$$

where A is a constant. If any p_i is assumed to be zero then $0 \ln 0$ is to be interpreted as zero. Since the constant A is present, logarithm can be taken to any base. Usually the logarithm is taken to the base 2 for ready application to binary systems. We will take logarithm to the base e .

1.3 Generalization of Shannon entropy

Consider again a multinomial population $P = (p_1, \dots, p_k)$, $p_i > 0, i = 1, \dots, k, p_1 + \dots + p_k = 1$. The following are some of the generalizations of Shannon entropy S_k .

$$R_{k,\alpha}(P) = \frac{\ln(\sum_{i=1}^k p_i^\alpha)}{1 - \alpha}, \quad \alpha \neq 1, \alpha > 0, \quad (4)$$

(Rényi entropy of order α of 1961)

$$H_{k,\alpha}(P) = \frac{\sum_{i=1}^k p_i^\alpha - 1}{2^{1-\alpha} - 1}, \quad \alpha \neq 1, \alpha > 0 \quad (5)$$

(Havrda-Charvát entropy of order α of 1967)

$$T_{k,\alpha}(P) = \frac{\sum_{i=1}^k p_i^\alpha - 1}{1 - \alpha}, \quad \alpha \neq 1, \alpha > 0 \quad (6)$$

(Tsallis entropy of 1988)

$$M_{k,\alpha}(P) = \frac{\sum_{i=1}^k p_i^{2-\alpha} - 1}{\alpha - 1}, \quad \alpha \neq 1, \quad -\infty < \alpha < 2 \quad (7)$$

(entropic form of order α)

$$M_{k,\alpha}^*(P) = \frac{\ln(\sum_{i=1}^k p_i^{2-\alpha})}{\alpha - 1}, \alpha \neq 1, \quad -\infty < \alpha < 2, \tag{8}$$

(additive entropic form of order α).

When $\alpha \rightarrow 1$ all the entropies of order α described above in (4) to (7) go to Shannon entropy S_k .

$$\lim_{\alpha \rightarrow 1} R_{k,\alpha}(P) = \lim_{\alpha \rightarrow 1} H_{k,\alpha}(P) = \lim_{\alpha \rightarrow 1} T_{k,\alpha}(P) = \lim_{\alpha \rightarrow 1} M_{k,\alpha}(P) = \lim_{\alpha \rightarrow 1} M_{k,\alpha}^*(P) = S_k. \tag{9}$$

Hence all the above measures are called generalized entropies of order α .

Let us examine to see what happens to the above entropies in the case of a joint distribution. Let $p_{ij} > 0, i = 1, \dots, m, j = 1, \dots, n$ such that $\sum_{i=1}^m \sum_{j=1}^n p_{ij} = 1$. This is a bivariate situation of a discrete distribution. Then the entropy in the joint distribution, for example,

$$M_{m,n,\alpha}(P, Q) = \frac{\sum_{i=1}^m \sum_{j=1}^n p_{ij}^{2-\alpha} - 1}{\alpha - 1}. \tag{10}$$

If the PPP holds and if $p_{ij} = p_i q_j, p_1 + \dots + p_m = 1, q_1 + \dots + q_n = 1, p_i > 0, i = 1, \dots, m, q_j > 0, j = 1, \dots, n$ and if $P = (p_1, \dots, p_m), Q = (q_1, \dots, q_n)$ then

$$\begin{aligned} (\alpha - 1)M_{m,\alpha}(P) + M_{n,\alpha}(Q) &= \frac{1}{\alpha - 1} \left(\sum_{i=1}^m p_i^{2-\alpha} - 1 \right) \left(\sum_{j=1}^n q_j^{2-\alpha} - 1 \right) \\ &= \frac{1}{\alpha - 1} \left[\sum_{i=1}^m \sum_{j=1}^n p_i^{2-\alpha} q_j^{2-\alpha} - \sum_{i=1}^m p_i^{2-\alpha} - \sum_{j=1}^n q_j^{2-\alpha} + 1 \right] \\ &= M_{m,n,\alpha}(P, Q) - M_{m,\alpha}(P) - M_{n,\alpha}(Q). \end{aligned}$$

Therefore

$$M_{m,n,\alpha}(P, Q) = M_{m,\alpha}(P) + M_{n,\alpha}(Q) + (\alpha - 1)M_{m,\alpha}(P)M_{n,\alpha}(Q). \tag{11}$$

If any one of the above mentioned generalized entropies in (4) to (8) is written as $F_{m,n,\alpha}(P, Q)$ then we have the relation

$$F_{m,n,\alpha}(P, Q) = F_{m,\alpha}(P) + F_{n,\alpha}(Q) + a(\alpha)F_{m,\alpha}(P)F_{n,\alpha}(Q). \tag{12}$$

where

$$\begin{aligned} a(\alpha) &= 0 \text{ (Rényi entropy } R_{k,\alpha}(P)) \\ &= 2^{1-\alpha} - 1 \text{ (Havrda-Charvát entropy } H_{k,\alpha}(P)) \\ &= 1 - \alpha \text{ (Tsallis entropy } T_{k,\alpha}(P)) \\ &= \alpha - 1 \text{ (entropic form of order } \alpha, \text{ i.e., } M_{k,\alpha}(P)) \\ &= 0 \text{ (additive entropic form of order } \alpha, \text{ i.e., } M_{k,\alpha}^*(P)). \end{aligned} \tag{13}$$

When $a(\alpha) = 0$ the entropy is called additive and when $a(\alpha) \neq 0$ the entropy is called non-additive. As can be expected, when a logarithmic function is involved, as in the cases of $S_k(P), R_{k,\alpha}(P), M_{k,\alpha}^*(P)$, the entropy is additive and $a(\alpha) = 0$.

1.4 Extensions to higher dimensional joint distributions

Consider a trivariate population or a trivariate discrete distribution $p_{ijk} > 0$, $i = 1, \dots, m$, $j = 1, \dots, n$, $k = 1, \dots, r$ such that $\sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^r p_{ijk} = 1$. If the PPP holds mutually, that is, pair-wise as well as jointly, which then will imply that

$$p_{ijk} = p_i q_j s_k, \quad \sum_{i=1}^m p_i = 1, \quad \sum_{j=1}^n q_j = 1, \quad \sum_{k=1}^r s_k = 1,$$

$$P = (p_1, \dots, p_m), \quad Q = (q_1, \dots, q_n), \quad S = (s_1, \dots, s_r).$$

Then proceeding as before, we have for any of the measures described above in (4) to (8), calling it $F(\cdot)$,

$$F_{m,n,r,\alpha}(P, Q, S) = F_{m,\alpha}(P) + F_{n,\alpha}(Q) + F_{r,\alpha}(S) + a(\alpha)[F_{m,\alpha}(P)F_{n,\alpha}(Q) + F_{m,\alpha}(P)F_{r,\alpha}(S) + F_{n,\alpha}(Q)F_{r,\alpha}(S)] + [a(\alpha)]^2 F_{m,\alpha}(P)F_{n,\alpha}(Q)F_{r,\alpha}(S) \tag{14}$$

where $a(\alpha)$ is the same as in (13). The same procedure can be extended to any multi-variable situation. If $a(\alpha) = 0$ we may call the entropy additive and if $a(\alpha) \neq 0$ then the entropy is non-additive.

1.5 Crucial recursivity postulate

Consider the multinomial population $P = (p_1, \dots, p_k)$, $p_i > 0$, $i = 1, \dots, k$, $p_1 + \dots + p_k = 1$. Let the entropy measure to be determined through appropriate postulates be denoted by $H_k(P) = H_k(p_1, \dots, p_k)$. For $k = 2$ let

$$f(x) = H_2(x, 1 - x), \quad 0 \leq x \leq 1 \text{ or } x \in [0, 1]. \tag{15}$$

If another parameter α is to be involved in $H_2(x, 1 - x)$ then we will denote $f(x)$ by $f_\alpha(x)$. From (5) to (7) it can be seen that the generalized entropies of order α of Havrda-Charvát (1967), Tsallis (1988, 2004) and Shannon (1948) entropy satisfy the functional equation

$$f_\alpha(x) + b_\alpha(x)f_\alpha\left(\frac{y}{1-x}\right) = f_\alpha(y) + b_\alpha(x)f\left(\frac{x}{1-y}\right) \tag{16}$$

for $x, y \in [0, 1)$ with $x + y \in [0, 1]$, with the boundary condition

$$f_\alpha(0) = f_\alpha(1) \tag{17}$$

where

$$\begin{aligned} b_\alpha(x) &= 1 - x \text{ (Shannon entropy } S_k(P)) \\ &= (1 - x)^\alpha \text{ (Havrda-Charvát entropy } H_{k,\alpha}(P)) \\ &= (1 - x)^\alpha \text{ (Tsallis entropy } T_{k,\alpha}(P)) \\ &= (1 - x)^{2-\alpha} \text{ (entropic form of order } \alpha, \text{ i.e., } M_{k,\alpha}(P)). \end{aligned} \tag{18}$$

Observe that the normalizing constant at $x = \frac{1}{2}$ is equal to 1 for $H_{k,\alpha}(P)$ and it is different for other entropies. Thus equations (6),(7),(8), with the appropriate normalizing constants $f_\alpha(\frac{1}{2})$, can give characterization theorems for the various entropy measures. The form of $b_\alpha(x)$ is coming from the crucial recursivity postulate, assumed as a desirable property for the measures.

1.6 Continuous analogues

In the continuous case let $f(x)$ be the density function of a real random variable x . Then the various entropy measures, corresponding to the ones in (4) to (8) are the following:

$$R_\alpha(f) = \frac{1}{1-\alpha} \ln \left[\int_{-\infty}^{\infty} [f(x)]^\alpha dx \right], \alpha \neq 1, \alpha > 0 \tag{19}$$

(Rényi entropy of order α)

$$H_\alpha(f) = \frac{1}{2^{1-\alpha} - 1} \left[\int_{-\infty}^{\infty} [f(x)]^\alpha dx - 1 \right], \alpha \neq 1, \alpha > 0 \tag{20}$$

(Havrda-Charvát entropy of order α)

$$T_\alpha(f) = \frac{1}{1-\alpha} \left[\int_{-\infty}^{\infty} [f(x)]^\alpha dx - 1 \right], \alpha \neq 1, \alpha > 0, \tag{21}$$

(Tsallis entropy of order α)

$$M_\alpha(f) = \frac{1}{\alpha-1} \left[\int_{-\infty}^{\infty} [f(x)]^{2-\alpha} dx - 1 \right], \alpha \neq 1, \alpha < 2 \tag{22}$$

(entropic form of order α)

$$M_\alpha^*(f) = \frac{1}{\alpha-1} \ln \left[\int_{-\infty}^{\infty} [f(x)]^{2-\alpha} dx \right], \alpha \neq 1, \alpha < 2 \tag{23}$$

(additive entropic form of order α).

As expected, Shannon entropy in this case is given by

$$S(f) = -A \int_{-\infty}^{\infty} f(x) \ln f(x) dx \tag{24}$$

where A is a constant.

Note that when PPP (product probability property) or statistical independence holds then in the continuous case also we have the property in (12) and (14) and then non-additivity holds for the measures analogous to the ones in (3),(5),(6),(7) with $a(\alpha)$ remaining the same. Since the steps are parallel a separate derivation is not given here.

2. Maximum entropy principle: distributional pathway

If we have a multinomial population $P = (p_1, \dots, p_k)$, $p_i > 0, i = 1, \dots, k, p_1 + \dots + p_k = 1$ or the scheme $P(A_i) = p_i, A_1 \cup \dots \cup A_k = S, P(S) = 1, A_i \cap A_j = \phi, i \neq j$ then we know that the maximum uncertainty in the scheme or the minimum information from the scheme is obtained when we cannot give any preference to the occurrence of any particular event or when the events are equally likely or when $p_1 = p_2 = \dots = p_k = \frac{1}{k}$. In this case, Shannon entropy becomes,

$$S_k(P) = S_k\left(\frac{1}{k}, \dots, \frac{1}{k}\right) = -A \sum_{i=1}^k \frac{1}{k} \ln \frac{1}{k} = A \ln k \quad (25)$$

and this is the maximum uncertainty or maximum Shannon entropy in this scheme. If the arbitrary functional f is to be fixed by maximizing the entropy then in (19) to (21) we have to optimize $\int_{-\infty}^{\infty} [f(x)]^\alpha dx$ for fixed α , over all functional f , subject to the condition $\int_{-\infty}^{\infty} f(x) dx = 1$ and $f(x) \geq 0$ for all x . For applying calculus of variation procedure we consider the functional

$$U = [f(x)]^\alpha - \lambda[f(x)]$$

where λ is a Lagrangian multiplier. Then the Euler equation is the following:

$$\frac{\partial U}{\partial f} = 0 \Rightarrow \alpha f^{\alpha-1} - \lambda = 0 \Rightarrow f = \left(\frac{\lambda}{\alpha}\right)^{\frac{1}{\alpha-1}} = \text{constant}. \quad (26)$$

Hence f is the uniform density in this case, analogous to the equally likely situation in the multinomial case. If the first moment $E(x) = \int_{-\infty}^{\infty} xf(x) dx$ is assumed to be a given quantity for all functional f then U will become the following for (19) to (21).

$$U = [f(x)]^\alpha - \lambda_1[f(x)] - \lambda_2 xf(x)$$

and the Euler equation leads to the power law. That is,

$$\frac{\partial U}{\partial f} = 0 \Rightarrow \alpha f^{\alpha-1} - \lambda_1 - \lambda_2 x = 0 \Rightarrow f = c_1 \left[1 + \frac{\lambda_2}{\lambda_1} x\right]^{\frac{1}{\alpha-1}}. \quad (27)$$

By selecting $c_1, \lambda_1, \lambda_2$ appropriately we can create a density out of (27). For $\alpha > 1$ and $\frac{\lambda_2}{\lambda_1} > 0$ the right side in (27) increases exponentially. If $\alpha = q > 1$ and $\frac{\lambda_2}{\lambda_1} = q - 1$ then we have Tsallis' q -exponential function from the right side of (27). If $\alpha > 1$ and $\frac{\lambda_2}{\lambda_1} = -(\alpha - 1)$ then (27) can produce a density in the category of a type-1 beta. From (27) it is seen that the form of the entropies of Havrda-Charvát $H_{k,\alpha}(P)$ and Tsallis $T_{k,\alpha}(P)$ need special attention to produce densities (Ferri et al. 2005). However, Tsallis has considered a different constraint on $E(x)$. If the density $f(x)$ is replaced by its escort density, namely, $\mu[f(x)]^\alpha$ where $\mu^{-1} = \int_x [f(x)]^\alpha dx$ and if the expected value of x in this escort density is assumed to be fixed for all functional f then the U of (26) becomes

$$U = f^\alpha - \lambda_1 f + \mu \lambda_2 x f^\alpha \quad \text{and}$$

$$\begin{aligned} \frac{\partial U}{\partial f} &= 0 \Rightarrow \alpha f^{\alpha-1}[1 + \mu\lambda_2 x] = \lambda_1 \Rightarrow f = \frac{\lambda^*}{(1 + \lambda_3 x)^{\frac{1}{\alpha-1}}} \Rightarrow \\ f &= \lambda_1^*[1 + \lambda_3 x]^{-\frac{1}{\alpha-1}} \end{aligned}$$

where λ_3 is a constant and λ_1^* is the normalizing constant. If λ_3 is taken as $\lambda_3 = \alpha - 1$ then

$$f = \lambda_1^*[1 + (\alpha - 1)x]^{-\frac{1}{\alpha-1}}. \tag{28}$$

Then (28) for $\alpha > 1$ is Tsallis statistics (Tsallis 2004; Cohen 2005). Then for $\alpha < 1$ also by writing $\alpha - 1 = -(1 - \alpha)$ one gets the case of Tsallis statistics for $\alpha < 1$ (Ferri et al. 2005). These modifications and the consideration of escort distribution are not necessary if we take the generalized entropy of order α . Thus if we consider $M_\alpha(f)$ and if we assume that the first moment in $f(x)$ itself is fixed for all functional f then the Euler equation gives

$$(2 - \alpha)f^{1-\alpha} - \lambda_1 + \lambda_2 x = 0 \Rightarrow f = \bar{\lambda} \left[1 - \frac{\lambda_2}{\lambda_1} x \right]^{\frac{1}{1-\alpha}}$$

and for $\frac{\lambda_2}{\lambda_1} = 1 - \alpha$ we have Tsallis statistics (Tsallis 2004, Cohen 2005)

$$f = \bar{\lambda}[1 - (1 - \alpha)x]^{\frac{1}{1-\alpha}} \tag{29}$$

coming directly, where $\bar{\lambda}$ is the normalizing constant.

Let us start with $M_\alpha(f)$ of (20) under the assumptions that $f(x) \geq 0$ for all x , $\int_a^b f(x)dx = 1$, $\int_a^b x^\delta f(x)dx$ is fixed for all functional f and for a specified $\delta > 0$, $f(a)$ is the same for all functional f , $f(b)$ is the same for all functional f , for some limits a and b , then the Euler equation becomes

$$(2 - \alpha)f^{1-\alpha} - \lambda_1 - \lambda_2 x^\delta = 0 \Rightarrow f = c_1[1 + c_1^* x^\delta]^{\frac{1}{1-\alpha}}. \tag{30}$$

If c_1^* is written as $-s(1 - \alpha)$, $s > 0$ then we have, writing f_1 for f ,

$$f_1 = c_1[1 - s(1 - \alpha)x^\delta]^{\frac{1}{1-\alpha}}, \delta > 0, \alpha < 1, 0 \leq x \leq \frac{1}{[s(1 - \alpha)]^{\frac{1}{\delta}}} \tag{31}$$

where $1 - s(1 - \alpha)x^\delta > 0$. For $\alpha < 1$ or $-\infty < \alpha < 1$ the right side of (31) remains as a generalized type-1 beta model with the corresponding normalizing constant c_1 . For $\alpha > 1$, writing $1 - \alpha = -(\alpha - 1)$ the model in (31) goes to a generalized type-2 beta form, namely,

$$f_2 = c_2[1 + s(\alpha - 1)x^\delta]^{-\frac{1}{\alpha-1}}. \tag{32}$$

When $\alpha \rightarrow 1$ in (31) or in (32) we have an extended or stretched exponential form,

$$f_3 = c_3 e^{-s x^\delta}. \tag{33}$$

If c_1^* in (30) is taken as positive then (30) for $\alpha < 1$, $\alpha > 1$, $\alpha \rightarrow 1$ will be increasing

exponentially. Hence all possible forms are available from (30). The model in (31) is a special case of the distributional pathway model and for a discussion of the matrix-variate pathway model see Mathai (2005). Special cases of (31) and (32) for $\delta = 1$ are Tsallis statistics (Gell-Mann & Tsallis 2004; Ferri et al. 2005).

Instead of optimizing $M_\alpha(f)$ of (22) under the conditions that $f(x) \geq 0$ for all x , $\int_a^b f(x)dx = 1$ and $\int_a^b x^\delta f(x)dx$ is fixed, let us optimize under the following conditions: $f(x) \geq 0$ for all x , $\int_a^b f(x)dx < \infty$ and the following two moment-like expressions are fixed quantities for all functional f ,

$$\int_a^b x^{(\gamma-1)(1-\alpha)} f(x)dx = \text{fixed}, \int_a^b x^{(\gamma-1)(1-\alpha)+\delta} f(x)dx = \text{fixed}.$$

Then the Euler equation becomes

$$(2-\alpha)f^{1-\alpha} - \lambda_1 x^{(\gamma-1)(1-\alpha)} - \lambda_2 x^{(\gamma-1)(1-\alpha)+\delta} = 0 \Rightarrow \\ f = c x^{\gamma-1} [1 + c^* x^\delta]^{\frac{1}{1-\alpha}}$$

and for $c^* = -s(1-\alpha)$, $s > 0$, we have the distributional pathway model for the real scalar case, namely

$$f(x) = c x^{\gamma-1} [1 - s(1-\alpha)x^\delta]^{\frac{1}{1-\alpha}}, \delta > 0, s > 0 \quad (34)$$

where c is the normalizing constant. For $\alpha < 1$, (34) gives a generalized type-1 beta form, for $\alpha > 1$ it gives a generalized type-2 beta form and for $\alpha \rightarrow 1$ we have a generalized gamma form. For $\alpha > 1$, (34) gives the superstatistics of Beck (2006) and Beck & Cohen (2003). For $\gamma = 1, \delta = 1$, (34) gives Tsallis statistics (Tsallis 2004; Cohen 2005). Densities appearing in a number of physical problems are seen to be special cases of (34), a discussion of which may be seen from Mathai & Haubold (2006a). For example, (34) for $\delta = 2, \gamma = 3, \alpha \rightarrow 1, x > 0$ is the Maxwell-Boltzmann density; for $\delta = 2, \gamma = 1, \alpha \rightarrow 1, -\infty < x < \infty$ is the Gaussian density; for $\gamma = \delta, \alpha \rightarrow 1$ is the Weibull density. For $\gamma = 1, \delta = 2, 1 < q < 3$ we have the Wigner function $W(p)$ giving the atomic moment distribution in the framework of Fokker-Planck equation, see Douglas, Bergamini, and Renzoni (2006) where

$$W(p) = z_q^{-1} [1 - \beta(1-q)p^2]^{\frac{1}{1-q}}, 1 < q < 3. \quad (35)$$

Before closing this section we may observe one more property for $M_\alpha(f)$. As an expected value

$$M_\alpha(f) = \frac{1}{\alpha-1} [E[f(x)]^{1-\alpha} - 1]. \quad (36)$$

But Kerridge's (Kerridge 1961) measure of "inaccuracy" in assigning $q(x)$ for the true density $f(x)$, in the generalized form is

$$H_\alpha(f : q) = \frac{1}{(2^{1-\alpha} - 1)} [E[q(x)]^{\alpha-1} - 1], \quad (37)$$

which is also connected to the measure of directed divergence between $q(x)$ and $f(x)$. In (37) the normalizing constant is $2^{1-\alpha} - 1$, the same factor appearing in Havrda-Charvát entropy. With different normalizing constants, as seen before, (36) and (37) have the same forms as an expected value with $q(x)$ replaced by $f(x)$ in (36). Hence $M_\alpha(f)$ can also be looked upon as a type of directed divergence or “inaccuracy” measure.

3. Associated differential pathway

The functional part in (34), for a more general exponent, namely

$$g(x) = \frac{f(x)}{c} = x^{\gamma-1} [1 - s(1 - \alpha)x^\delta]^{\frac{\beta}{1-\alpha}}, \quad \alpha \neq 1, \delta > 0, \beta > 0, s > 0 \tag{38}$$

is seen to satisfy the following differential equation for $\gamma \neq 1$ which defines the differential pathway.

$$x \frac{d}{dx} g(x) = (\gamma - 1)x^{\gamma-1} [1 - s(1 - \alpha)x^\delta]^{\frac{\beta}{1-\alpha}} - s\beta\delta x^{\delta+\gamma-1} [1 - s(1 - \alpha)x^\delta]^{\frac{\beta}{1-\alpha} [1 - \frac{(1-\alpha)}{\beta}]}. \tag{39}$$

Then for $\delta = \frac{(\gamma-1)(\alpha-1)}{\beta}$, $\gamma \neq 1$, $\alpha > 1$ we have

$$x \frac{d}{dx} g(x) = (\gamma - 1)g(x) - s\beta\delta [g(x)]^{1 - \frac{(1-\alpha)}{\beta}} \tag{40}$$

$$= (\gamma - 1)g(x) - s\delta [g(x)]^\alpha \tag{41}$$

for $\beta = 1, \gamma \neq 1, \delta = (\gamma - 1)(\alpha - 1), \alpha > 1$.

For $\gamma = 1, \delta = 1$ in (38) we have

$$\frac{d}{dx} g(x) = -s[g(x)]^\eta, \quad \eta = 1 - \frac{(1 - \alpha)}{\beta} \tag{42}$$

$$= -s[g(x)]^\alpha \text{ for } \beta = 1. \tag{43}$$

Here (39) is the differential equation for the diverse distribution functions generated through the entropic pathway (Souza & Tsallis 2005) and (43) is the power law coming from Tsallis statistics (Gell-Mann & Tsallis 2004).

Remark. The referee initiated the attention of the authors to the *Editorial for the proceedings of the NEXT2003 conference on news and expectations in thermostatics* by G. Kaniadakis and M. Lissia, available at arXiv:cond-mat/0409615v1. The authors join the spirit of Kaniadakis and Lissia that parameter interpretation in information theory and mathematical statistics is less important and that Boltzmann-Gibbs-Shannon entropy has been generalized in different ways by adding one or even two parameters. This paper provides the mathematical and statistical methods to unify the generalizations of such entropies in the case of one parameter through the entropic and distributional pathways. It also supplements the more diversified distribution functions with a differential pathway.

Acknowledgments

The authors express their appreciation for enlightening conversations with Professor Constantino Tsallis, Centro Brasileiro de Pesquisas Físicas Brazil and Santa Fe Institute USA, on the topic of this paper at the occasion of UN/ESA/NASA workshops on basic space science. The authors also would like to thank the Department of Science and Technology, Government of India, New Delhi, for the financial assistance for this work under project No. SR/S4/MS:287/05 which enabled this collaboration possible.

References

- Beck, C., 2006 *Physica A*, 365, 96-101
 Beck, C., & Cohen, E.G.D., 2003, *Physica A*, 322, 267
 Cohen, E.G.D., 2005, *Pramana*, 64, 635
 Douglas, P., Bergamini, S., & Renzoni, F., 2006, *PhRvL*, 96, 110601-1-4
 Ferri, G.L., Martinez, S., & Plastino, A., 2005, *JSMTE*, PO4009
 Gell-Mann, M. & Tsallis, C. (eds) 2004, *Nonextensive Statistical Mechanics: Interdisciplinary Applications*, Oxford University Press, Oxford
 Havrda, J., & Charvát, F., 1967, *Kybernetika*, 3, 30
 Kerridge, D.F., 1961, *J. Royal Statistical Society Series B*, 23, 184
 Mathai, A.M., 2005, *Linear Algebra and Its Applications*, 396, 317
 Mathai, A.M., & Haubold, H.J., 2006, *Physica A*, 375, 110
 Mathai, A.M., & Haubold, H.J., 2006a, On generalized distributions and pathways, arXiv:cond-mat/0609526v2
 Mathai, A.M., & Rathie, P.N., 1975, *Basic Concepts in Information Theory and Statistics: Axiomatic Foundations and Applications*, Wiley Halstead: New York and Wiley Eastern, New Delhi
 Rényi, A., 1961, *Proc. Fourth Berkeley Symposium on Mathematical Statistics and Probability*, 1960, University of California Press, 1961, Vol. 1, 547
 Shannon, C.E., 1948, *Bell System Technical Journal*, 27, 379, 547
 Souza, A.M., & Tsallis, C., 2005, *Proc. 31st Workshop of the International School of Solid State Physics 20-26 July 2004, Erice, Italy*, eds C. Beck et al., World Scientific, Singapore, 66 [arXiv:cond-mat/0501389v1]
 Tsallis, C., 1988, *JSP*, 52, 479
 Tsallis, C., 2004, *Physica D*, 193, 3