

Galaxies: Structure, formation and evolution

Lecture 20



Yogesh Wadadekar

Mar-Apr 2022

Question


At a given stellar mass, how do star formation, morphology, and environment of galaxies depend on one another?

Dependence of star-formation on morphology




Bait et al. (2017)

Dependence of star-formation on morphology




Bait et al. (2017)

Dependence of morphology on environment




Bait et al. (2017)

Dependence of star-formation on environment




Bait et al. (2017)

Galactic fountain model makes hot halo



Classical and Pseudo bulges



Classical and Pseudo bulges

Classical

steep Sérsic index

Pseudo

shallow Sérsic index

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals

Pseudo

shallow Sérsic index
common in LT spirals

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals
old stellar population

Pseudo

shallow Sérsic index
common in LT spirals
Mix of old/young stars

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals
old stellar population
formed like elliptical galaxies

Pseudo

shallow Sérsic index
common in LT spirals
Mix of old/young stars
secular evolution

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals
old stellar population
formed like elliptical galaxies
stellar bar absent

Pseudo

shallow Sérsic index
common in LT spirals
Mix of old/young stars
secular evolution
stellar bar important

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals
old stellar population
formed like elliptical galaxies
stellar bar absent
follow Kormendy relation and FP

Pseudo

shallow Sérsic index
common in LT spirals
Mix of old/young stars
secular evolution
stellar bar important
do not follow relation

Classical and Pseudo bulges

Classical

steep Sérsic index
common in ET Spirals
old stellar population
formed like elliptical galaxies
stellar bar absent
follow Kormendy relation and FP
in high mass galaxies

Pseudo

shallow Sérsic index
common in LT spirals
Mix of old/young stars
secular evolution
stellar bar important
do not follow relation
in low mass galaxies

How does pseudo/classical occurrence depend on environment? Are their stellar ages correlated with the disc? Why are classical bulges more common in S0s?

Mishra et al. 2017a, 2017b, 2018

Formation mechanisms for ellipticals and spirals

- Eggen, Lynden-Bell & Sandage (1962) considered a model in which galaxies form from the collapse of gas clouds, and suggested that the difference between ellipticals and spirals reflects the rapidity of star formation during the collapse.
- If most of the gas turns into stars as it falls in, the collapse is effectively dissipationless and infall motions are converted into the random motion of stars, resulting in an elliptical.
- If the cloud remains gaseous during collapse, the gravitational energy can be effectively dissipated via shocks and radiative cooling. In this case, the cloud will shrink until it is supported by angular momentum, leading to the formation of a rotationally supported disk.

Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars

Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes which are much more complicated than LSS formation and growth

Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes which are much more complicated than LSS formation and growth
- Probably closely related to the formation of the massive central black holes as well

Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes which are much more complicated than LSS formation and growth
- Probably closely related to the formation of the massive central black holes as well
- Generally, we think of massive galaxy formation at high redshifts ($z \sim 3 - 10$, say); dwarfs may be still forming now (**cosmic downsizing**)

Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes which are much more complicated than LSS formation and growth
- Probably closely related to the formation of the massive central black holes as well
- Generally, we think of massive galaxy formation at high redshifts ($z \sim 3 - 10$, say); dwarfs may be still forming now (**cosmic downsizing**)
- Observations have found populations of what must be young galaxies (ages < 1 Gyr), ostensibly progenitors of large galaxies today, at $z \sim 5 - 7$


Galaxy formation

- has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes which are much more complicated than LSS formation and growth
- Probably closely related to the formation of the massive central black holes as well
- Generally, we think of massive galaxy formation at high redshifts ($z \sim 3 - 10$, say); dwarfs may be still forming now (**cosmic downsizing**)
- Observations have found populations of what must be young galaxies (ages < 1 Gyr), ostensibly progenitors of large galaxies today, at $z \sim 5 - 7$
- The JWST frontier is at $z \sim 7 - 20$, the so-called Reionization Era

Galaxy formation timeline from the star formation perspective


- The smallest scale density fluctuations keep collapsing, with baryons falling into the potential wells dominated by the dark matter, achieving high densities through cooling
- This process starts right after the recombination at $z \sim 1100$
- Once the gas densities are high enough, star formation ignites probably around $z \sim 20 - 30$
- By $z \sim 6$, UV radiation from young galaxies reionizes the universe
- These protogalactic fragments (of stars and gas) keep merging, forming larger objects in a hierarchical fashion. During this,
- Star formation enriches the gas, and some of it is expelled in the intergalactic medium, while more gas keeps falling in
- If a central massive black hole forms, the energy release from accretion can also create considerable feedback

Dark matter mass function evolves too




Barkana and Loeb (2001)

Halo, subhaloes, centrals and (missing) satellites




Can baryonic properties be determined by the properties of the dark matter halo? See: Paranjape et al. (2018)

Halo merger trees- hierachical “bottom up” growth



Galaxy formation simulations match reality




Why is galaxy formation difficult?

Galaxy formation is actually a much messier problem than structure formation. In addition to gravity and build-up of host dark halos (fairly well understood) we need to add relatively poorly understood baryonic processes:

- Shock heating of gas
- Cooling of gas into dark halos
- Formation of stars (also not a well understood process!) from the cold gas
- The evolution of the resulting stellar population
- Feedback processes generated by the ejection of mass and energy from evolving stars
- Production and mixing of heavy elements (chemical evolution)
- Effects of dust obscuration
- Formation of black holes at galaxy centers and effects of AGN emission, jets, etc.


Evolution of an individual galaxy



Mo et al. (2010)


Galaxy formation recipe

A Recipe for Galaxy Formation




Baugh, Cole, Frenk & Lacey

Galaxy formation schematic




Galaxy formation flowchart - simplified reality



Mo et al. (2010)

Galaxy formation and evolution in one diagram



see review by Kormendy & Kennicutt (2004)